



US Army Corps
of Engineers

ENVIRONMENTAL IMPACT RESEARCH PROGRAM

MISCELLANEOUS PAPER EL-86-

APPLICATION OF THE HABITAT EVALUATION

PROCEDURES IN THE CYPRESS BAYOU

BASIN, TEXAS

by

K. Jack Killgore

Environmental Laboratory

DEPARTMENT OF THE ARMY

Waterways Experiment Station, Corps of Engineers

PO Box 631


Vicksburg, Mississippi 39180-0631

and

Paul M. Hathorn

US Army Engineer District, Fort Worth

Fort Worth, Texas 76102-0300



Final Report

December 1986

Approved For Public Release; Distribution Unlimited

Prepared for US Army Engineer District, Fort Worth
Fort Worth, Texas 76102-0300

PRELIMINARY



REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188 Exp. Date Jun 30 1986	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS PRELIMINARY DRAFT		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Miscellaneous Paper EL-86-			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION USAEWES Environmental Laboratory		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) PO Box 631 Vicksburg, MS 39180-0631			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION USAED, Fort Worth		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) 19 Taylor St. Fort Worth, Texas 76102-0300			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Application of the Habitat Evaluation Procedures in the Cypress Bayou Basin, Texas					
12. PERSONAL AUTHOR(S) Killgore, K. Jack, Hathorn, Paul M.					
13a. TYPE OF REPORT Final report		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) December 1986	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION Available from National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Compensation Habitat Evaluation Lake		
			Fishes Impact River		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Construction of a dam on either the Little or Black Cypress Bayou, Texas, is being evaluated by the US Army Engineer District, Fort Worth, to provide water resource benefits (flood control, water supply, recreation) in the Cypress Bayou Basin. The Habitat Evaluation Procedure (HEP) was used to determine fish habitat gains in the proposed reservoirs, estimate losses in fish habitat resulting from inundation of portions of the Little and Black Cypress bayous, and recommend methods to compensate for habitat losses caused by the project. Habitat Suitability Index (HSI) models composed of three physical habitat variables (depth, velocity, and cover) were developed from field and literature data for nine species of riverine fishes including: spotted bass, longear sunfish, grass pickerel, flathead catfish, blacktail shiner, ironcolor shiner, brook silverside, spotted sucker, and slough (Continued)					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

Preface

PRELIMINARY DRAFT

This report describes an aquatic resource evaluation of a proposed water resource project in the Cypress Bayou Basin, Texas, and contributes to the overall feasibility study being prepared by the US Army Engineer District, Fort Worth (SWF). Funding for this project was provided by SWF; partial funding for development of the Suitability Index Curves was provided by the Environmental Impact Research Program (Work Unit 32390).

The study was completed by the Aquatic Habitat Group (AHG), Environmental Resources Division (ERD), Environmental Laboratory (EL), US Army Engineer Waterways Experiment Station (WES). The report was prepared by Messrs. K. Jack Killgore (AHG) and Paul M. Hathorn (SWF). Mr. Tom Cloud (US Fish and Wildlife Service, Fort Worth), Mr. Mike Ryan (Texas Parks and Wildlife Department), Dr. Andrew Miller (WES), Dr. William Matthews (University of Oklahoma), Mr. Kenneth Conley (WES), and Mr. Frank Ferguson (WES) contributed to the conduct of this study. The report was prepared under the supervision of Dr. Thomas Wright, Chief, AHG; Dr. Conrad J. Kirby, Chief, ERD; and Dr. John Harrison, Chief, EL. This report was edited by Ms. Lee T. Byrne of the WES Information Technology Laboratory.

COL Allen F. Grum, USA, was the previous Director of WES. COL Dwayne G. Lee, CE, is the present Commander and Director. Dr. Robert W. Whalin is Technical Director.

This report should be cited as follows:

Killgore, K. J., and Hathorn, P. M. 1986. "Application of the Habitat Evaluation Procedure in the Cypress Bayou Basin, Texas," Miscellaneous Paper EL-86- , US Army Engineer Waterways Experiment Station, Vicksburg, Miss.

Contents

PRELIMINARY DRAFT
Page

Preface	1
List of Tables	3
List of Figures	3
Conversion Factors, Non-SI to SI (Metric) Units of Measurement	4
Introduction	5
Purpose and Objectives	6
Methods	6
Study area	6
River models	7
Lake models	9
Field methods--rivers	9
Data analysis	11
Results	12
Discussion	19
Conclusions and Recommendations	21
References	22
Appendix A: Summary of Water Quality Variables in the Cypress Bayou Basin, Texas	A1
Appendix B: Flow Duration Tables for the Little and Black Cypress Bayous, Texas	B1
Appendix C: Fish Species List of the Little and Black Cypress Bayous, Texas	C1
Appendix D: Feeding and Reproductive Guild of Fishes in the Cypress Bayou Basin	D1
Appendix E: Monthly Periodicity of Evaluation Species Relative to Temperature and Discharge	E1
Appendix F: Suitability Index Curves for the Nine Evaluation Riverine Species	F1
Appendix G: Description of Hydraulic Analysis to Predict Physical Habitat for Unmeasured Flows	G1
Appendix H: Composite Habitat Unit Discharge Table for the Little and Black Cypress Bayous	H1
Appendix I: Habitat Unit Duration Table for the Cypress Bayou Basin HEP Study	I1
Appendix J: Habitat Units Lost from Inundation in the Little Cypress Bayou	J1

List of Tables

PRELIMINARY DRAFT

<u>No.</u>		<u>Page</u>
1	Overview of the Steps Taken to Conduct an Aquatic HEP for the Cypress Bayou Basin Project	5
2	Delineation Between the River and Lake Habitat for the Little and Black Cypress Bayous	8
3	Summary of Regression Equations and Variables Used to Calculate HSI Values for Lake Evaluation Species	10
4	Maintenance Flows for the Little and Black Cypress Bayous	18
5	Compensation Flows for the Little Cypress Bayou	19
6	Average Annual Habitat Units (HU's) of Lake Species for Marshall and Black Cypress Lake During the Time Period of 1 to 10 Years	20

List of Figures

1	Location of study sites in the Cypress Bayou Basin, Texas	7
2	Schematic drawing of procedure to estimate habitat availability for unmeasured flows	12
3	Habitat unit (HU) discharge plots of the evaluation species for the Little Cypress Bayou	13
4	Habitat unit (HU) discharge plots of the evaluation species for the Black Cypress Bayou	14
5	Plots of the habitat unit (HU) duration values and flow for the Little Cypress Bayou	16
6	Plots of the habitat unit (HU) duration values and flow for the Black Cypress Bayou	17

PRELIMINARY DRAFT

Conversion Factors, Non-SI to SI (Metric)
Units of Measurement

Non-SI units of measurement used in this report can be converted to SI (metric) units as follows:

<u>Multiply</u>	<u>By</u>	<u>To Obtain</u>
acres	4,046.873	square metres
cubic feet	0.02831685	cubic metres
feet	0.3048	metres
miles (US statute)	1.609347	kilometres
square feet	0.09290304	square metres

Application of the Habitat Evaluation Procedure
in the Cypress Bayou Basin, Texas

PRELIMINARY DRAFT

Introduction

1. The US Army Engineer District, Fort Worth (SWF), is investigating the feasibility of providing flood control, water supply, recreation, and other water resource benefits for the Cypress Bayou Basin, located in north-east Texas. Of the alternative plans considered, construction of a dam on either the Little Cypress Bayou (Marshall Lake) or Black Cypress Bayou (Black Cypress Lake) appears to be the most feasible approach to accommodate the various water resource needs in the basin. Aquatic resource studies of the project were initiated in 1984 by a team of biologists representing SWF, US Fish and Wildlife Service (USFWS), Texas Parks and Wildlife Department (TPWD), and Waterways Experiment Station (WES). The Habitat Evaluation Procedure (HEP) was selected as one method to evaluate the impacts of the project on aquatic resources. The study approach follows the format described in the HEP manual (USFWS 1980) with modifications specific-to-project requirements. An overview of the steps taken in the HEP analysis appears in Table 1.

Table 1
Overview of the Steps Taken to Conduct an Aquatic HEP
for the Cypress Bayou Basin Project

-
- | | |
|---------|---|
| Step 1: | Delineate the river and future lake habitat and describe the hydraulic and morphometric features. |
| Step 2: | Select evaluation fish species and construct the Habitat Suitability Index (HSI) models. |
| Step 3: | Select representative reaches, collect hydraulic and morphometric data, and estimate physical habitat conditions at target discharges using hydraulic mathematical relationships. |
| Step 4: | Construct habitat duration curves and define maintenance flows. |
| Step 5: | Determine habitat units lost in the river due to inundation and develop a plan to compensate for lost habitat. |
| Step 6: | Determine habitat gains of the project created by the reservoirs. |
-

Purpose and Objectives

2. The purpose of this document is to provide SWF with a comprehensive analysis of fish habitat gains and losses resulting from the construction of a dam on either Little or Black Cypress Bayou. The objectives are:

- a. To determine baseline habitat conditions that would maintain the historic fish community structure.
- b. To recommend techniques to compensate for the loss of inundated fish habitat.
- c. To identify gains in new fish habitat created by the reservoir.

MethodsStudy area

3. The study area included the Little and Black Cypress bayous located in northeastern Texas (Figure 1). Both rivers are lowland, meandering, warm-water streams that are relatively undisturbed by water resource development. The rivers have abundant instream cover such as logjams, rootwads, undercut banks, and cypress trees. Substrate composition is relatively uniform ranging from clayey sand to silty clay. Based on data from the US Geological Survey (USGS) gaging stations located on both rivers near Jefferson, Texas, water quality (Appendix A) is adequate to sustain viable fish populations at any flow and therefore was not used in the HEP analysis. The average annual discharge for the Little and Black Cypress bayous is 527 and 333 cfs*, respectively. Discharge ranges from 0 during August through October to greater than 1,000 cfs during the spring months (Appendix B).

4. Three major study areas were used in the HEP: the rivers below the damsites, the lakes, and the portion of rivers that would be inundated (Table 2). The river habitats below the dams extend from the damsite downstream to the confluence with the Big Cypress Creek. The river reaches that would be inundated by the project are between the damsite and the conservation pool elevation (US Army Engineer District, Fort Worth (SWF) 1985).

* A table of factors for converting non-SI units of measurement to SI (metric) units is presented on page 4.




Figure 1. Location of study sites in the Cypress Bayou Basin, Texas

River models

5. From a total of 67 species of fishes known to occur in both rivers (Appendix C), nine evaluation species were chosen for the HEP. These were spotted bass, grass/chain pickerel, flathead catfish, longear sunfish, spotted sucker, blacktail shiner, ironcolor shiner, brook silverside, and slough darter. These species were selected from biological guilds (Appendix D) that considered adult feeding preferences and reproductive strategies and represented 87 percent of the fish community. All evaluation species were considered to be equally important to the stream ecosystem. A periodicity table (Appendix E) was constructed to relate the presence of life stages (spawning, fry, juvenile, and adults) to changes in discharge and water temperature.

Table 2
Delineation Between the River and Lake Habitat for the
Little and Black Cypress Bayous

<u>River</u>	<u>Type of Habitat</u>	<u>River Channel miles</u>	<u>Elevation ft</u>	<u>acres</u>
Little Cypress	River below damsite	1-20.3	170-195	646*
	Inundated river	20.3-41.3	195-255	132**
	Lake (conservation pool)	--	195-255	28,988
Black Cypress	River below damsite	1-17.0	175-200	194*
	Inundated river	17-44.0	200-253	--
	Lake (conservation pool)	--	200-253	21,951

* Calculated at annual median flow occurring at USGS Gage near Jefferson, Texas.

** Calculated at annual median flow occurring at USGS Gage near Ore City, Texas.

6. The variables used to assess fish habitat were depth, velocity, and cover. These physical habitat variables are important because they: (a) regulate the carrying capacity of a river system if water quality is within the tolerance limits of the species, (b) are directly impacted by water resource development, (c) can be manipulated to provide optimum habitat conditions, and (d) are easily measured in the field. Suitability Index (SI) Curves for these variables were developed from field data for all evaluation species except the flathead catfish and slough darter. Curves for these species were developed from the literature. Because of the lack of field data on nonadult life stages, only adult life stages were used in the HSI models. Juveniles generally occurred in habitats similar to those of adults. Requirements for spawning and for survival of fry were accounted for by the occurrence of overbank flows.

7. Fish habitat utilization was determined by measuring water depth, water velocity, and the presence or absence of instream cover at each location where an evaluation species was captured by electrofishing. Length and weight

of each evaluation species were recorded at the time of capture to separate the species into adults, juveniles, and fry. To the extent possible, an equal amount of time was spent at each type of habitat (channel, side channel, and shoreline). Field data were collected seasonally for both rivers during 1984.

8. SI curves were prepared for each evaluation species (Appendix F). These curves summarize the frequency of capture for each of the three habitat variables and for each evaluation species. The Y-axis, or SI Score, ranges from 0.0 (poor habitat) to 1.0 (optimum habitat) and is a qualitative measure of habitat value. An average HSI score for each species was derived from the geometric mean of all variables using the following formula:

$$HSI = (V_1 \cdot V_2 \cdot V_3)^{0.333} \quad (1)$$

where

HSI = Habitat Suitability Index value for physical habitat

V_1 = depth, ft

V_2 = velocity, ft/sec

V_3 = cover, percent

Lake models

9. The following fishes were evaluated for the proposed lakes: large-mouth bass, bluegill, black crappie, white bass, total sport fishes, and total fishes. Predicted standing crops for each species were determined using regression equations prepared by the USFWS (Table 3) and were converted to HSI scores using the technique described in Aggus and Morais (1979).

Field methods--rivers

10. Prior to field sampling, a reconnaissance of both rivers was made by boat, and two representative sites were selected at each river. The sites on the Little Cypress Bayou were located at river mile 2 (Elevation 170 ft, represented 13 river miles) and near the Highway 154 Bridge crossing (Elevation 210 ft, represented 7.3 river miles). Sites on the Black Cypress Bayou were located at river miles 1.5 (Elevation 175 ft, represented 10.5 river miles) and near Berea Bridge crossing (Elevation 200 ft, represented 6.5 river miles). At each site, a metal tag line was positioned across the river at two locations separated by 0.1 mile, and depth, velocity, and

Table 3

Summary of Regression Equations and Variables Used to Calculate

HSI Values for Lake Evaluation Species

Species	Regression Equation to Predict Standing Crop	R ²	Lake Habitat Variables							Age years
			Lake	Dissolved Solids mg/l	Growing Season days	Outlet Depth ft	Area of Conservation Pool ft ²	Mean Depth ft	Water Level Fluctuation ft	
Largemouth bass	0.5743 - 0.3120 (log water level fluctuations) + 0.2594 (log dissolved solids) + 0.0046 (age)	0.244	Little Cypress (Marshall Lake)	150	213	60	28,988	23	10	1
Bluegill	-821.4815 + 366.5507 (log growing season) - 0.0688 (dissolved solids) + 0.00006 (dissolved solids squared)	0.244	Black Cypress	50	213	60	21,951	20	10	1
Black crappie	2.7778 - 0.0088 (dissolved solids) - 0.00001 (dissolved solids squared)	0.500								
White bass	5.1756 - 0.9512 (log area) - 2.9939 (log mean depth) + 0.0309 (outlet depth) + 1.2550 (log dissolved solids)	0.172								
Total sport fish	0.9809 - 0.0056 (mean depth) + 0.3877 (log mean depth) + 0.9944 (log growing season)	0.094								
Total species	4.9397 + 0.1614 (log area) - 0.0090 (mean depth) - 1.2663 (log growing season) - 2 x 10 ⁻⁸ (dissolved solids squared)	0.292								

PRELIMINARY DRAFT

cover were measured at regular intervals (number of intervals = 10 percent of the cross-sectional width) that divided the cross section into cells. Water depth was measured to the nearest 0.1 ft using a leveling rod. Water velocity was measured to the nearest 0.1 ft/sec using a Marsh-McBirney model 201 current meter. If the total depth (TD) was less than or equal to 3.0 ft, then velocity was measured at 0.6 TD. If TD exceeded 3.0 ft, then velocity was measured at both 0.2 and 0.8 TD, and an average was obtained. Cover was classified as "present" or "not present" in each cell and converted into the percentage of cells with cover. In addition, the slope and distance from the water's edge to the high-water mark were measured with a hand-held level and tape measure respectively. Collectively, these sites represented habitat below the damsites. In addition, the downstream transect site at Highway 154 represented habitat features above the damsite for the Little Cypress Bayou.

Data analysis

11. A noncomputerized method of determining depth, velocity, cover, and other morphometric features of the cross sections at a range of discharges, partially modified from Dunham and Collotzi (1975) and Bovee and Milhous (1978), was used to predict physical habitat conditions at unmeasured flows. The water surface profile measured in the field was plotted on graph paper (Figure 2), and unmeasured hydraulic geometric features of the cross sections were extracted from these graphs in order to calculate velocity and to determine the water depth and percentage of cover for a range of discharges. A detailed description of this procedure for the Little Cypress Bayou is shown in Appendix G.

12. HU's were determined from the following equation:

$$HU = HSI \times \text{Acres} \quad (2)$$

where

HSI = Habitat Suitability Index

Acres = Acres of river at a given discharge

HU = Habitat units

This equation was applied to each discharge of interest (10 to 1,000 cfs) for each species at each representative reach. An SI was assigned to the value of each variable (depth, velocity, cover) that occurred at the target discharges.




Figure 2. Schematic drawing of procedure to estimate habitat availability for unmeasured flows

The SI values were aggregated into the HSI model to obtain a value between 0.0 to 1.0 that indicated the suitability of the conditions of depth, velocity, and cover to the evaluation species. The product of the HSI equation was multiplied by the acres of river that occur at each target discharge to obtain HU's. (Total HU's) for the river were calculated by adding the HU's of the representative reaches for each target discharge.

Results

13. An increase in discharge usually resulted in a positive change in HU's for all species (Figures 3 and 4). HU's increased most rapidly between 0 and 200 cfs, and either tapered off or slightly decreased at discharges greater than 200 cfs. Decreases in HU's were due to high velocities without any substantial addition of cover. HU's increased at overbank flows (i.e., 425 and 460 cfs for the Little and Black Cypress bayous, respectively) because of an increase in cover, shallow depths, and surface area. The Little Cypress Bayou provided more fish habitat than the Black Cypress Bayou provided at all





Figure 3. Habitat unit (HU) discharge plots of the evaluation species for the Little Cypress Bayou

PRELIMINARY DRAFT





Figure 4. Habitat unit (HU) discharge plots of the evaluation species for the Black Cypress Bayou

Same FWS, add
stop further
work to complete
the study.
FWS
stop further
work to complete
the study.

discharges. Species that preferred or could tolerate high-velocity, deep water (such as the blacktail shiner, spotted bass, spotted sucker, and flathead catfish) had higher amounts of HU's than did species that usually inhabited shallow, slow-moving water with substantial amounts of instream cover (such as the pickerel, longear sunfish, brook silverside, ironcolor shiner and slough darter). Even though the amounts of HU's were different among species for a given discharge, the trend of the HU discharge curves was similar. Therefore, to simplify data interpretation, a composite HU discharge curve was developed from the average of all nine individual species curves and was used to determine baseline conditions and compensation requirements (Figures 3 and 4).

* 14. Maintenance flows have been defined for this study as the positive, inflection point on an HU duration-discharge curve and are considered to be those baseline conditions that would maintain the historic fish community structure for a specific time period. An HU duration curve is a cumulative frequency plot that shows the percentage of a certain amount of habitat being equalled or exceeded during a given time period, as described in Bovee (1982). A 10-percent value indicates HU's that occur infrequently, whereas a 90-percent value indicates HU's that occur frequently. For each river, the 10-through 90-percent HU duration values were plotted on the y axis, and the flows that corresponded with each HU value were indicated on the x axis (Figures 5 and 6). The inflection points were visually interpreted from these figures and from a table of these data (Appendix H). Table 4 shows the monthly maintenance flows for each river. The maintenance flows for most months occurred around the 60-percent HU exceedance value. maintenance flows during the late winter and spring ranged from 190 to 270 cfs in both rivers and declined to near 0 cfs in the summer and early fall.

15. The Little and Black Cypress bayous are classified by USFWS as resource category 2 (in-kind replacement, no trade-offs); therefore, habitat gains from the lake were not included in the compensation analysis. Due to a determination late in the study that a damsite on Black Cypress Bayou was not economically feasible, a compensation plan was conducted for only the Little Cypress Bayou. Loss in HU's at the 50-percent exceedance flow was determined by month to represent the portion of the Little Cypress lost as the result of inundation. The monthly 50-percent exceedance flows were obtained from the

PRELIMINARY DRAFT




Figure 5. Plots of the habitat unit (HU) duration values and flow for the Little Cypress Bayou. (Arrows indicate the inflection point corresponding to the maintenance flow)

10-90% flows → Computed habitat -
Exp. > 100 cfs — 0 cfs } from fig. 4 values -

high flows
flow duration from 10 to 90% streamflow
flow duration uses from 10 to 90% streamflow




Figure 6. Plots of the habitat unit (HU) duration values and flow for the Black Cypress Bayou. (Arrows indicate the inflection point corresponding to the maintenance flow)

Table 4

Maintenance Flows for the Little and Black Cypress Bayous

PRELIMINARY DRAFT

Month	Maintenance Flow, cfs	
	Little Cypress Bayou	Black Cypress Bayou
January	190	190
February	215	210
March	215	270
April	270	210
May	270	180
June	40	55
July	14	7
August	3	3
September	3	3
October	3	3
November	16	65
December	55	180

USGS gaging station at Highway 259 near Ore City, because it more accurately represented the flows occurring in the overall river segment that would be inundated than did the downstream gaging station (i.e., Highway 59). Furthermore, HSI values and other morphometric features, including acres, that occurred at each median monthly discharge at the USGS gage near Ore City were determined from the Highway 154 downstream transect (see Table G3), which was considered representative of the inundated stream habitat of the Little Cypress Bayou. The total HU's lost to lake habitat ranged from 333 to 1,502 depending upon the season (Appendix J). It was determined that compensation flows of 10 to greater than 425 cfs (i.e., overbank flows) would be needed below the dam to achieve full and in-kind compensation for habitat lost to inundation (Table 5) and to maintain the historic fish community from the dam-site to the mouth of the Little Cypress Bayou.

16. An aquatic HEP was conducted for the proposed Marshall and Black Cypress lakes (Table 6). The analysis includes a 10-year period beginning immediately after dam closure and assumes that the physical and chemical variables used in the lake HSI models (Table 3) would not significantly change

cf. this out. I think they would change.

Table 5

Compensation Flows for the Little Cypress Bayou

PRELIMINARY DRAFT

Month	Maintenance Flow Below the Dam		Monthly Median Flow at USGS Gage near Ore City	Habitat Units Lost from Inundation	Compensation Flow	
	cfs	HU's			cfs	HU's
January	190	2,420	149	1,011	>425*	>3,000
February	215	2,500	253	1,448	>425	>3,000
March	215	2,500	298	1,502	>425	>3,000
April	270	2,600	206	1,212	>425	>3,000
May	270	2,600	193	876	>425	>3,000
June	40	1,010	45	487	100	1,500
July	14	850	6	314	50	1,160
August	3	400	2	333	10	700
September	3	400	2	333	10	700
October	3	400	3	333	10	700
November	16	990	33	442	85	1,400
December	55	1,110	92	760	150	1,900

sum = $\frac{1294}{12}$

$$\bar{M} = 107 \text{ cfs} = 77,465 \text{ ac-ft/yr.} = 60.1\% \text{ of firm annual yield}$$

* Overbank flows.

during this time period. Marshall Lake had the highest amount of habitats for all species except bluegill. These data were prepared to define habitat gains from the project and were not intended to facilitate trade-off analysis. With either lake, however, these gains would occur and should be considered as intangible benefits of the lake. These values can also be used in determination of economic man-days (recreation) benefits attributable to the lake project.

Discussion

17. Rivers in the Cypress Bayou Basin undergo extreme seasonal water level fluctuations. Summer drought accompanied by high-water temperatures and low dissolved oxygen (see Appendix A) drastically decreases usable fish habitat. These conditions can increase spatial competition for food and habitat

Table 6

Average Annual Habitat Units (HU's) of Lake Species for Marshall and
Black Cypress Lake During the Time Period of 1970 to 1980

DRAFT

Lake	Species	Area of Habitat acres	Habitat Suitability Index	Average Annual HU's
Marshall Lake (Little Cypress)	All Species	28,988	0.75	21,741
	Bluegill	28,988	0.45	13,045
	Largemouth Bass	28,988	0.40	11,595
	Black Crappie	28,988	0.50	14,494
	White Bass	28,988	0.78	22,610
	Sportfish	28,988	0.58	16,813
Black Cypress Lake	All Species	21,951	0.77	16,902
	Bluegill	21,951	0.71	15,585
	Largemouth Bass	21,951	0.35	7,683
	Black Crappie	21,951	0.62	13,609
	White Bass	21,951	0.65	14,268
	Sportfish	21,951	0.55	12,073

(Cowx, Young, and Hellawell 1984) and can also increase foraging efficiency by predators because of clear water and concentrated prey (Stevens and Miller 1983). In contrast, high flows during spring increase usable fish habitat and ensure adequate spawning, survival, and nursery habitat for fishes. Instream flow releases, particularly during the summer drought, would moderate standing crop fluctuations in downstream reaches and compensate for in-kind habitat lost from inundation.

18. The HEP is a flexible procedure to assess changes in habitat from water resource projects. A variety of species-oriented assessment techniques have been developed that are conceptually similar to HEP but differ in expertise (training) requirements, time and resource constraints, data requirements, and objectives pursued (Schuytema 1982, Coulombe 1978). The HEP is ideally suited for analyzing lake habitat, although limited by one's ability to predict future habitat conditions. This method is specifically tailored to

PRELIMINARY DRAFT

facilitate trade-off analysis and to develop compensation plans. The HEP was selected to analyze river habitat to minimize the requirements for data acquisition and analysis as well as to provide a quantitative and relatively rapid approach in determining changes in fish habitat as a function of flow. An important advantage in using the hydraulic procedures described in this report was the ability to extrapolate the amount of usable fish habitat to a flow range of 0 to 1,000 cfs in a relatively short time. Six working days were required to complete the river analysis, including the collection of field data (physical habitat), and to determine maintenance plus compensation flows.

Conclusions and Recommendations

19. Usable habitat for nine species of fish increased with discharge up to 200 cfs, moderated or decreased at flows from 200 to 400 cfs, and again increased at overbank flows.

20. The longear sunfish, ironcolor shiner, grass/chain pickerel, and slough darter preferred shallow, slow-moving water with abundant instream cover, whereas the spotted bass, blacktail shiner, spotted sucker, and flat-head catfish liked deeper water with moderate to fast flow usually associated with large instream objects such as cypress trees and logjams. The brook silverside was found in both types of habitat.

21. To maintain the status quo of the fish community structure below the proposed damsite, the monthly maintenance flows that appear in Table 5 should be released.

22. To compensate for the inundated fish habitat, the compensation flows that appear in Table 6 should be released. Overbank flows should be released periodically during the spring spawning season to maximize spawning areas and to ensure fry survival.

23. Marshall Lake will create more fish habitat than will Black Cypress Lake.

Aggus, L. R., and Morais, D. I. 1979. "Habitat Suitability Index Equations for Reservoirs Based on Standing Crop of Fish," US Fish and Wildlife Service, Fort Collins, Colo.

Bovee, K. D. 1982. "A Guide to Stream Habitat Analysis Using the Instream Flow Incremental Methodology," Instream Flow Information Paper No. 12, FWS/OBS-82/26, US Fish and Wildlife Service, Fort Collins, Colo.

Bovee, K. D., and Milhous, R. 1978. "Hydraulic Simulation in Instream Flow Studies: Theory and Techniques," Instream Flow Information Paper: No. 5, FWS/OBS-78/33, US Fish and Wildlife Service, Fort Collins, Colo.

Coulombe, H. N. 1978 (Jan.). "Toward an Integrated Ecological Assessment of Wildlife Habitat," National Workshop of Integrated Inventories of Renewable National Resources, Tucson, Ariz.

Cowx, I. G., Young, W. O., and Hellawell, J. M. 1984. "The Influence of Drought on the Fish and Invertebrate Populations of an Upland Stream in Wales," Freshwater Biology, Vol 14, pp 165-177.

Dunham, D. K., and Collotzi, A. 1975. "The Transect Method of Stream Habitat Inventory," US Forest Service, Ogden, Utah.

Grossman, G. D. 1982. "Dynamics and Organization of a Rocky Intertidal Fish Assemblage: the Persistence and Resilience of Taxocene Structure," American Naturalist, Vol 119, pp 611-637.

Henderson, F. M. 1966. Open Channel Flow, MacMillan Co., New York.

Ross, R. T., Matthews, W. J., and Echelle, A. A. 1985. "Persistence of Stream Fish Assemblages: Effects of Environmental Change," The American Naturalist, Vol 126, No. 1, pp 24-40.

Schuytema, G. S. 1982. "A Review of Aquatic Habitat Assessment Methods," EPA-600/3-82-002, US Environmental Protection Agency, Corvallis, Oreg.

Stevens, D. E., and Miller, L. E. 1983. "Effects of River Flow on Abundance of Young Chinook Salmon, American Shad, Longfin Smelt, and Delta Smelt in the Sacramento-San Joaquin River System," North American Journal of Fisheries Management, Vol 3, pp 425-437.

US Army Engineer District, Fort Worth (SWF). 1985. "Cypress Bayou Basin Feasibility Study-Formulation," Fort Worth, Texas.

US Fish and Wildlife Service (USFWS). 1980. "Habitat Evaluation Procedures," ESM 102, Fort Collins, Colo.

Appendix A: Summary of Water Quality Variables in the Cypress Bayou Basin, Texas

River/Month	Conductivity $\mu\text{mho/cm}$	Temperature $^{\circ}\text{C}$	Total		Dissolved Oxygen mg/l	Turbidity JTU
			Dissolved Solids mg/l			
<u>Little Cypress Bayou</u>						
January	178 \pm 58(11)	5.9 \pm 2.6(11)	110 \pm 30(11)	10.9 \pm 0.8(7)	8.0 \pm 5.0(4)	
February	186 \pm 54(6)	8.0 \pm 3.4(6)	110 \pm 25(6)	8.7 \pm 1.7(3)	--	
March	153 \pm 43(9)	15.4 \pm 1.7(9)	94 \pm 25(9)	7.2 \pm 1.1(5)	6.0 \pm 1.7(3)	
April	146 \pm 56(8)	17.7 \pm 1.8(8)	86 \pm 30(8)	6.9 \pm 0.9(6)	15.0 \pm 2.8(2)	
May	185 \pm 99(8)	22.2 \pm 1.5(7)	111 \pm 56(8)	6.1 \pm 0.3(4)	26.0 \pm 1.4(2)	
June	150 \pm 76(7)	24.7 \pm 2.4(7)	93 \pm 45(7)	5.5 \pm 1.3(7)	14.0 \pm 12.7(2)	
July	207 \pm 59(8)	27.5 \pm 1.5(8)	125 \pm 26(8)	4.3 \pm 1.1(5)	23.5 \pm 10.6(2)	
August	296 \pm 206(9)	27.1 \pm 2.3(9)	213 \pm 148(9)	5.2 \pm 0.6(5)	24.0 \pm 25.4(2)	
September	182 \pm 55(8)	24.5 \pm 2.9(7)	109 \pm 29(8)	5.4 \pm 0.7(4)	14.0(1)	
October	220 \pm 82(9)	17.5 \pm 3.0(9)	127 \pm 47(9)	6.9 \pm 1.6(5)	5.0(1)	
November	225 \pm 91(9)	11.3 \pm 4.0(9)	133 \pm 48(9)	8.8 \pm 2.4(4)	7.5 \pm 0.7(2)	
December	189 \pm 104(9)	8.5 \pm 2.1(9)	109 \pm 54(9)	9.9 \pm 2.0(4)	--	
<u>Black Cypress Bayou</u>						
January	52 \pm 4(5)	7.6 \pm 1.1(4)	45 \pm 4(5)	--	--	
February	56 \pm 11(7)	8.2 \pm 4.3(7)	45 \pm 8(7)	--	--	
March	57 \pm 15(4)	16.6 \pm 0.7(4)	42 \pm 5(4)	--	--	
April	56 \pm 12(6)	16.1 \pm 4.3(6)	43 \pm 9(6)	--	--	
May	53 \pm 11(6)	20.2 \pm 1.7(6)	43 \pm 10(6)	--	--	
June	62 \pm 15(8)	25.1 \pm 2.5(8)	48 \pm 10(8)	--	--	
July	69 \pm 14(7)	27.4 \pm 1.8(7)	50 \pm 5(7)	--	--	
August	86 \pm 34(6)	26.8 \pm 2.4(6)	58 \pm 15(6)	--	--	
September	86 \pm 41(3)	28.3 \pm 3.5(3)	63 \pm 19(3)	--	--	
October	63 \pm 8(6)	18.4 \pm 4.1(6)	50 \pm 4(6)	--	--	
November	82 \pm 62(7)	10.2 \pm 3.4(7)	59 \pm 28(7)	--	--	
December	59 \pm 18(5)	7.7 \pm 3.8(5)	46 \pm 14(4)	--	--	

PRELIMINARY

PRELIMINARY DRAFT

Appendix B: Flow Duration Table for the Little and Black Cypress Bayous, Texas

Site/Percent Exceedance	Discharge by Month, cfs											
	January	February	March	April	May	June	July	August	September	October	November	December
	Little Cypress Bayou*											
10	1,560	1,750	1,950	2,070	2,430	1,090	352	100	226	251	801	1,360
20	1,010	1,300	1,430	1,320	1,520	618	158	43	84	111	300	853
30	782	997	1,060	941	1,030	315	95	27	38	49	159	511
40	581	807	870	702	720	194	54	17	21	31	104	317
50	409	658	689	555	531	135	30	9	9	15	72	227
60	277	543	546	413	371	89	18	4	2	6	47	163
70	205	407	411	282	238	54	11	1	0	1	23	112
80	120	245	289	192	135	30	4	0	0	0	8	72
90	70	131	192	113	85	14	1	0	0	0	0	45
n	1,178	1,074	1,178	1,140	1,178	1,170	1,209	1,209	1,170	1,178	1,140	1,178
	Black Cypress Bayou*											
10	919	952	1,390	1,560	1,010	737	147	74	221	220	530	962
20	696	760	956	919	675	391	75	29	27	86	292	636
30	531	645	761	614	473	245	49	16	12	44	187	457
40	399	546	632	456	336	167	35	9	6	21	117	324
50	307	465	510	340	244	121	21	5	2	11	78	231
60	236	384	394	256	179	72	8	2	0	3	49	169
70	182	312	321	208	138	40	3	1	0	1	31	133
80	194	237	253	159	99	16	1	0	0	0	13	100
90	111	161	187	102	51	4	0	0	0	0	4	75
n	496	452	496	480	496	480	496	496	480	527	480	496

PRELIMINARY DRAFT

* Flow calculated from USGS gages at Highway 59 near Jefferson, Texas.

Appendix C: Fish Species List of the Little and Black Cypress

Bayous, Texas

Checklist of Fish Species Collected from the Little and Black
Cypress Rivers, Texas. Collected by Ryan, Matthews, Killgore
(1984) - O; collected by Kemp (1954) - X; not collected - NC

PRELIMINARY DRAFT

Common Name	Species	Little Cypress	Black Cypress
Chestnut lamprey	<i>Ichthyomyzon castaneus</i>	X	NC
Spotted gar	<i>Lepisosteus oculatus</i>	O	NC
Longnose gar	<i>Lepisosteus osseus</i>	X	NC
Bowfin	<i>Amia calva</i>	O	O
Gizzard shad	<i>Dorosoma cepedianum</i>	O	O
Grass pickerel	<i>Esox americanus</i> <i>vermiculatus</i>	O	O
Chain pickerel	<i>Esox niger</i>	O	O
Black buffalo	<i>Ictiobus niger</i>	X	NC
Smallmouth buffalo	<i>Ictiobus bubalus</i>	X	NC
Spotted sucker	<i>Minytrema melanops</i>	O	O
Common carp	<i>Cyprinus carpio</i>	O	O
Golden shiner	<i>Notemigonus crysoleucas</i>	O	X
Pugnose minnow	<i>Notropis emiliae</i>	O	O
Emerald shiner	<i>Notropis atherinoides</i>	O	X
Ribbon shiner	<i>Notropis fumeus</i>	O	O
Redfin shiner	<i>Notropis umbratilis</i>	O	O
Ironcolor shiner	<i>Notropis chalybaeus</i>	O	O
Weed shiner	<i>Notropis texanus</i>	O	O
Pallid shiner	<i>Notropis amnis</i>	O	O
Blacktail shiner	<i>Notropis venustus</i>	O	O
Red shiner	<i>Notropis lutrensis</i>	X	NC
Sand shiner	<i>Notropis stramineus</i>	X	NC
Blackspot shiner	<i>Notropis atrocaudalis</i>	X	X
Silvery minnow	<i>Hybognathus nuchalis</i>	X	X
Cypress minnow	<i>Hybognathus hayi</i>	X	X

(Continued)

(Sheet 1 of 3)

(Continued)

Common Name	Species	Little	Black
		Cypress	Cypress
Bullhead minnow	<i>Pimephales vigilax</i>	X	0
Channel catfish	<i>Ictalurus punctatus</i>	0	0
Black bullhead	<i>Ictalurus melas</i>	0	X
Yellow bullhead	<i>Ictalurus natalis</i>	0	X
Flathead catfish	<i>Pylodictis olivaris</i>	0	0
Tadpole madtom	<i>Noturus gyrinus</i>	0	0
American eel	<i>Anguilla rostrata</i>	0	0
Golden topminnow	<i>Fundulus chrysotus</i>	0	X
Starhead topminnow	<i>Fundulus blairae</i>	0	X
Blackstripe topminnow	<i>Fundulus notatus</i>	0	X
Blackspotted topminnow	<i>Fundulus olivaceus</i>	0	0
Mosquitofish	<i>Gambusia affinis</i>	0	0
Pirate perch	<i>Aphredoderus sayanus</i>	0	0
Brook silversides	<i>Labidesthes sicculus</i>	0	0
White bass	<i>Morone chrysops</i>	0	0
Yellow bass	<i>Morone mississippiensis</i>	0	NC
Spotted bass	<i>Micropterus punctulatus</i>	0	0
Largemouth bass	<i>Micropterus salmoides</i>	0	0
Warmouth	<i>Lepomis gulosus</i>	0	0
Green sunfish	<i>Lepomis cyanellus</i>	0	NC
Spotted sunfish	<i>Lepomis punctatus</i>	0	0
Bantam sunfish	<i>Lepomis symmetricus</i>	NC	X
Redear sunfish	<i>Lepomis microlophus</i>	0	0
Bluegill	<i>Lepomis macrochirus</i>	0	0
Orangespotted sunfish	<i>Lepomis humilis</i>	NC	X
Redbreast sunfish	<i>Lepomis auritus</i>	NC	X
Longear sunfish	<i>Lepomis megalotis</i>	0	0
Dollar sunfish	<i>Lepomis marginatus</i>	X	X
White crappie	<i>Pomoxis annularis</i>	0	0
Black crappie	<i>Pomoxis nigromaculatus</i>	0	0

(Continued)

(Sheet 2 of 3)

(Concluded)


Common Name	Species	Little Cypress	Black Cypress
Flier	<i>Centrarchus macropterus</i>	NC	0
Banded pygmy sunfish	<i>Elassoma zonatum</i>	0	X
Black side darter	<i>Percina maculata</i>	0	0
Dusky darter	<i>Percina sciera</i>	NC	X
Log perch	<i>Percina caprodes</i>	NC	0
Scaly sand darter	<i>Ammocrypta vivax</i>	NC	X
Bluntnose darter	<i>Etheostoma chlorosomum</i>	0	0
Slough darter	<i>Etheostoma gracile</i>	0	X
Mud darter	<i>Etheostoma asprigene</i>	0	NC
Cypress darter	<i>Etheostoma proeliare</i>	0	0
Redfin darter	<i>Etheostoma whipplei</i>	0	NC
Freshwater drum	<i>Aplodinotus grunniens</i>	0	0
Totals	67 species	60	56

(Sheet 3 of 3)

PRELIMINARY DRAFT

D1

Appendix E: Monthly Periodicity of Evaluation Species Relative to Temperature and Discharge



PRELIMINARY DRAFT

Appendix F: Suitability Index Curves for the Nine Evaluation
Riverine Fish Species

PRELIMINARY DRAFT




Figure F1. Suitability Index Curves for spotted bass and
spotted sucker adults

PRELIMINARY DRAFT




Figure F2. Suitability Index Curves for pickerel and flathead catfish adults

PRELIMINARY DRAFT




Figure F3. Suitability Index Curves for brook silverside and blacktail shiner adults

1, .85, .1

= .4/3

= .085

PRELIMINARY DRAFT




Figure F4. Suitability Index Curves for ironcolor shiner and longear sunfish adults

PRELIMINARY DRAFT




Figure F5. Suitability Index Curves for slough darter

*

Appendix G: Description of Hydraulic Analysis to Predict
Physical Habitat at Unmeasured Flows

PRELIMINARY DRAFT

1. The purpose of this appendix is to describe the procedure to determine the value of the physical habitat variables (depth, velocity, and cover) and other morphometric features for unmeasured flow conditions in the Little Cypress Bayou. These data are used to calculate HU's to determine maintenance and compensation flows.

2. The first step is to estimate the coefficient of roughness (n) and calculate the slope of the channel (Se) using field data. These values remain constant and are used to determine velocity for unmeasured flows. The coefficient of roughness ranges from 0.025 for clear and straight river channels to 0.150 for weedy and overgrown channels (Bovee and Milhous 1978, Henderson 1966). The coefficient of roughness used in the Little and Black Cypress bayous was 0.075. Once n has been estimated, the slope is calculated from the following equation:

$$Se = \frac{n^2 V^2}{2.22 R^{4/3}} \quad (G1)$$

where

V = mean channel velocity measured in the field, ft/sec

n = coefficient of roughness

$$R = \text{Hydraulic Radius} = \frac{\text{Area, ft}^2}{\text{Wetted Perimeter, ft}}$$

The values to calculate hydraulic radius (area and wetted perimeter) are determined from the graphs (Figure 2). Velocity is then calculated for each cell using Manning's equation expressed as follows:

$$V, \text{ ft/sec} = \frac{1.486}{n} R^{2/3} Se^{1/2} \quad (G2)$$

The calculated velocities are compared with the field-measured velocities to check the accuracy of the variables used in Manning's equation. If the

Manning No.

velocities do not agree, the slope is adjusted. In most cases, either no or small changes in the slope were required for this study. Once the cell velocities were determined, cell discharge was expressed as follows:

$$Q = V \cdot A$$

(G3)

where

Q = discharge, cfs

V = velocity, ft/sec

A = area, ft²

The cell discharges were summed to obtain a channel discharge that corresponded to the stage height on the graph (Figure 2).

3. Tables G1 and G2 illustrate the steps to determine depth, velocity, and acres of river that occur at the target discharges for the two representative study sites in the Little Cypress Bayou. Target discharges correspond to an incremental range of flows that could be released from the dam. The first step was to calculate the average depth, velocity, and width for each transect at discharges ranging from extreme low flows to overbank flows, using the hydraulic equations and graphs described in the previous paragraph. To accomplish this, new stage heights were drawn on the graph paper (Figure 2). From these graphs, the unmeasured hydraulic components (hydraulic radius and velocity) were determined. Discharge was also calculated for each new stage height. The second step was to calculate regression equations to predict the average depth, velocity, and width for a given discharge. The regression equations were then used to predict average depth, velocity, and width at target discharges of 10, 50, 100, 200, 300, 400, 500, and 1,000 cfs. For cover, a plot was made that related the percentage of cover (i.e., percentage of cells with cover) and discharge for each cross section. An average percentage of cover at each target discharge was then tabulated for each river. These data provided a depth, velocity, and percent cover at each discharge and at each representative site that was used to determine the HSI value. The fourth step was to determine the acres of river that occurred at each discharge by multiplying width times river miles. The final step was to calculate HU's for the study area at each target discharge using the method described in paragraph 12 (Table G3).

Table G1

Procedure to Determine Average Depths, Velocities, and Channel Widths over a Range of Flows Using the principle geometry information from the traps. Field Data Was Collected from the Little Cypress Bayou at Hwy 134

Step 1: Calculate average depth, velocity, and width for each transect at 4 discharges.

Downstream Transect				Upstream Transect			
Discharge cfs	Channel Width ft	Depth, ft $\bar{x} \pm SD(n)$	Velocity ft/sec $\bar{x} \pm SD(n)$	Discharge cfs	Channel Width ft	Depth, ft $\bar{x} \pm SD(n)$	Velocity ft/sec $\bar{x} \pm SD(n)$
20	51	1.0 \pm 0.61(6)	0.30 \pm 0.13(6)	9	41	1.8 \pm 0.62(4)	0.11 \pm 0.03(4)
81	66	2.2 \pm 0.92(7)	0.49 \pm 0.15(7)	81	60	5.3 \pm 1.90(6)	0.23 \pm 0.08(6)
232	93	3.1 \pm 1.70(10)	0.62 \pm 0.26(10)	200	93	6.6 \pm 3.90(9)	0.25 \pm 0.13(9)
449	230	3.7 \pm 2.20(13)	0.69 \pm 0.30(13)	556	430	6.8 \pm 5.10(20)	0.27 \pm 0.14(20)

Step 2: Calculate regression equations to predict the average depth, velocity, and width for a given discharge.

Downstream Transect			
Depth, ft = $Q (0.006) + 1.39$	$R^2 = 0.86$		
Velocity, ft/sec = $Q (0.0008) + 0.37$	$R^2 = 0.82$		
Width, ft = $Q (0.36) + 29.3$	$R^2 = 0.93$		

Upstream Transect			
Depth, ft = $Q (0.007) + 3.7$	$R^2 = 0.51$		
Velocity, ft/sec = $Q (0.0002) + 0.17$	$R^2 = 0.52$		
Width, ft = $Q (0.74) - 0.06$	$R^2 = 0.96$		

Step 3: Using the regression equations, calculate the average depth, velocity, and width between the upstream and downstream transects over the discharges of interest. Plot percent cover and discharge for each transect and take the average.

Discharge cfs	Depth, ft			Velocity, ft/sec			Width, ft			Cover percent
	Downstream + Upstream + 2 = Average Transect	Transect	Transect	Downstream + Upstream + 2 = Average Transect	Transect	Transect	Downstream + Upstream + 2 = Average Transect	Transect	Transect	
10	1.4	3.7	2.6	0.37	0.17	0.27	33	30	32	20
50	1.7	4.0	2.8	0.41	0.18	0.29	50	37	43	25
100	2.0	4.4	3.2	0.45	0.19	0.32	71	74	73	33
200	2.5	5.0	3.7	0.53	0.21	0.37	112	148	130	52
300	3.1	5.7	4.4	0.61	0.23	0.42	153	222	187	60
400	3.7	6.4	5.1	0.69	0.25	0.47	194	296	245	75
500	4.2	7.1	5.6	0.77	0.28	0.52	236	370	303	70
1000	7.1	10.5	8.8	1.20	0.38	0.79	442	740	591	90

Step 4: Calculate the acres of river that the two transects represent over the discharges of interest. This site represents 7.3 river miles. Use the following equation to obtain acres: Acres, ft² = [Width * (miles * 5,280)] * (2.296 * 10⁻⁵).

Discharge, cfs	Acres, ft ²
10	28
50	38
100	66
200	115
300	165
400	217
500	268
1,000	523

these are acres

conversion
Lower eq ft to acre.
0.0002296

$$32 \times (7.3 \times 5,280) \times (2.296 \times 10^{-5}) = 28.32$$

$$32 \times (1.6 \times 5,280) = 2.38$$

Table A2

Procedure to Determine Average Depths, Velocities, and Channel Widths over a Range of Flows Using the Hydraulic Geometry
Information from the Graphs. Field Data Was Collected from the Little Cypress Stream, 3.000 ft Wide.

PRELIMINARY DRAFT

Step 1: Calculate average depth, velocity, and width for each transect at 4 discharges.

Downstream Transect				Upstream Transect (Approximately 500 ft Upstream of Downstream Transect)			
Discharge cfs	Channel Width ft	Depth, ft $\bar{x} \pm SD(n)$	Velocity ft/sec $\bar{x} \pm SD(n)$	Discharge cfs	Channel Width ft	Depth, ft $\bar{x} \pm SD(n)$	Velocity ft/sec $\bar{x} \pm SD(n)$
11	28	1.8 \pm 1.1(3)	0.16 \pm 0.07(3)	8	96	1.1 \pm 0.45(8)	0.07 \pm 0.02(8)
44	64	3.0 \pm 1.9(6)	0.19 \pm 0.08(6)	82	123	3.8 \pm 1.2(12)	0.17 \pm 0.04(12)
141	124	3.7 \pm 2.3(12)	0.23 \pm 0.09(12)	225	147	5.8 \pm 2.3(15)	0.22 \pm 0.06(15)
289	350	4.3 \pm 3.0(19)	0.25 \pm 0.13(19)	398	250	7.0 \pm 3.3(19)	0.25 \pm 0.09(19)

Step 2: Calculate regression equations to predict the average depth, velocity, and width for a given discharge.

Downstream Transect			Upstream Transect		
Depth, ft = $Q (0.008) + 2.25$	$R^2 = 0.82$		Depth, ft = $Q (0.014) + 1.90$	$R^2 = 0.89$	
Velocity, ft/sec = $Q (0.0003) + 0.17$	$R^2 = 0.88$		Velocity, ft/sec = $Q (0.0004) + 0.103$	$R^2 = 0.82$	
Width, ft = $Q (1.14) + 3.41$	$R^2 = 0.96$		Width, ft = $Q (0.40) + 80.5$	$R^2 = 0.95$	

Step 3: Using the regression equations, calculate the average depth, velocity, and width between the upstream and downstream transects over the discharges of interest. Plot the percentage of cover and discharge for each transect and take the average.

Discharge cfs	Depth, ft			Velocity, ft/sec			Width, ft			Cover percent
	Downstream Transect	Upstream Transect	$\div 2 =$ Average	Downstream Transect	Upstream Transect	$\div 2 =$ Average	Downstream Transect	Upstream Transect	$\div 2 =$ Average	
10	2.3	2.0	2.15	0.17	0.10	0.14	15	84	50	35
50	2.6	2.6	2.60	0.18	0.12	0.16	60	100	80	40
100	3.0	3.3	3.15	0.20	0.14	0.17	117	120	119	50
200	3.8	4.7	4.25	0.23	0.19	0.21	231	160	196	60
300	4.6	5.1	5.40	0.26	0.23	0.25	345	200	272	68
400	5.4	5.6	6.50	0.29	0.27	0.28	459	240	349	73
500	6.2	9.0	7.60	0.32	0.31	0.32	573	280	426	78
10,000	10.1	16.1	13.00	0.47	0.52	0.50	1,142	479	810	80

Step 4: Calculate the acres of river that the two transects represent over the discharges of interest. This site represents 13 river miles.

Discharge, cfs	Acres, ft ²
10	79
50	126
100	187
200	309
300	429
400	550
500	671
1,000	1,276

FWS Maintenance Flow Need Recommendations:

75 - M - N (5 mo.)

100 - D - A (7 mo.)

Table G3
Habitat Suitability Index Values and Habitat Units for the Evaluation Species
in the Little Cypress Bayou

PRELIMINARY DRAFT

Site/Discharge	Total Acres Each Reach	Spotted Bass		Spotted Sucker		Brook Silverside		Blacktail Shiner		Ironcolor Shiner		Longear Sunfish		Pickereel		Flathead Catfish		Slough Carter		
		HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	HSI	HU's	
Little Cypress	Mile 2*																			
	10	79	0.76	60	0.62	49	0.80	63	0.65	51	0.76	60	0.89	70	0.89	71	0.68	54	0.74	58
	50	126	0.86	108	0.75	94	0.82	103	0.70	88	0.62	78	0.77	97	0.89	113	0.71	89	0.68	86
	100	187	0.90	168	0.93	168	0.89	166	0.76	142	0.56	105	0.66	123	0.86	161	0.75	140	0.60	112
	200	309	0.93	287	1.0	309	0.54	167	0.81	250	0.38	117	0.59	182	0.68	210	0.57	210	0.50	154
	300	429	0.65	279	0.60	257	0.35	150	0.81	347	0.28	124	0.38	163	0.33	142	0.61	262	0.42	180
	400	550	0.62	341	0.36	193	0.28	154	0.76	418	0.27	137	0.34	187	0.27	137	0.49	269	0.33	181
	500	671	0.62	362	0.34	228	0.22	148	0.48	322	0.26	174	0.34	228	0.24	168	0.38	255	0.34	228
	1,000	1,276	0.31	395	0.26	332	0.17	217	0.39	485	0.20	268	0.22	281	0.16	204	0.21	268	0.20	255
Little Cypress	Hwy 154**																			
	10	28	0.67	19	0.46	12	0.45	13	0.52	15	0.38	11	0.45	13	0.48	13	0.38	11	0.38	11
	50	38	0.74	28	0.53	20	0.53	20	0.58	22	0.38	13	0.43	16	0.52	20	0.38	14	0.42	16
	100	66	0.79	52	0.65	43	0.49	32	0.65	43	0.32	21	0.46	30	0.58	38	0.43	28	0.46	30
	200	115	0.89	102	0.79	91	0.50	57	0.80	92	0.33	38	0.50	57	0.55	63	0.45	45	0.33	38
	300	165	0.86	142	0.69	114	0.33	54	0.86	142	0.30	49	0.39	64	0.41	68	0.38	63	0.35	58
	400	217	0.63	137	0.50	108	0.29	63	0.96	208	0.32	69	0.27	59	0.26	56	0.38	82	0.20	54
	500	268	0.49	131	0.40	107	0.19	51	0.90	241	0.21	56	0.25	67	0.23	62	0.38	102	0.21	56
	1,000	523	0.23	120	0.17	89	0.17	89	0.39	204	0.20	105	0.18	94	0.16	84	0.26	136	0.20	105
Little Cypress	Damsite to mouth†																			
	10	107	--	79	--	61	--	76	--	66	--	71	--	83	--	84	--	65	--	69
	50	164	--	136	--	114	--	123	--	110	--	91	--	113	--	133	--	103	--	102
	100	253	--	220	--	204	--	198	--	185	--	126	--	153	--	199	--	168	--	142
	200	424	--	389	--	400	--	224	--	342	--	155	--	239	--	273	--	255	--	192
	300	646	--	421	--	371	--	204	--	489	--	173	--	227	--	210	--	325	--	238
	400	767	--	478	--	306	--	217	--	626	--	206	--	246	--	193	--	351	--	237
	500	939	--	493	--	335	--	199	--	563	--	230	--	295	--	230	--	357	--	284
	1,000	1,799	--	515	--	421	--	306	--	689	--	373	--	375	--	288	--	404	--	360

Damsite = Hwy 154 (upstream site) → RM 2 (downstream site)

* Represents 13 river miles.
** Represents 7.3 river miles.
† Represents 20.3 miles.

Appendix H: Composite Habitat Unit--Discharge Table
for the Little and Black Cypress Bayous

PRELIMINARY DRAFT

Discharge cfs	Habitat Units	
	<u>Little Cypress Bayou</u>	<u>Black Cypress Bayou</u>
0	300	200
10	654	440
50	1,025	575
100	1,595	759
200	2,469	986
300	2,658	1,154
400	2,860	1,213
500	2,986	1,326
1,000	3,730	1,699

Appendix I: Habitat Unit Duration Table for the Cypress Bayou Basin HEP Study

Month/ Site	Habitat Unit Duration Values, percent*																	
	10%		20%		30%		40%		50%		60%		70%		80%		90%	
	HU	Flow	HU	Flow	HU	Flow	HU	Flow	HU	Flow	HU	Flow	HU	Flow	HU	Flow	HU	Flow
Little Cypress: Mile 1-Dam site																		
January	3,670	900	3,375	670	3,060	530	2,850	390	2,655	290	2,420	190	2,035	155	1,500	90	1,020	45
February-March	4,080	1,200	3,740	1,050	3,450	740	3,260	640	2,965	485	2,810	360	2,650	285	2,500	215	2,000	155
April-May	4,300	2,000	3,775	1,075	3,380	670	3,000	510	2,795	370	2,600	270	2,200	175	1,500	90	1,025	50
June	3,500	760	2,990	510	2,500	215	2,000	155	1,600	105	1,300	75	1,011	40	805	12	540	8
July	2,575	265	1,730	110	1,250	60	1,000	30	850	14	565	7	520	6	350	2	300	0
August-October	2,000	155	1,275	65	810	13	725	11	654	10	400	3	350	2	325	1	300	0
November	3,000	510	2,500	215	1,775	125	1,400	85	1,250	60	990	16	800	12	600	8	375	3
December	3,690	930	3,175	610	2,820	375	2,420	190	2,180	170	1,805	130	1,450	88	1,110	55	950	15
Black Cypress: Mile 0-Dam site																		
January	1,900	1,300	1,705	1,050	1,510	730	1,315	480	1,120	280	975	190	910	170	856	140	800	120
February-March	1,940	1,325	1,800	1,200	1,660	940	1,520	740	1,379	570	1,238	420	1,100	270	975	190	880	145
April	2,500	2,000	1,847	1,250	1,629	900	1,410	600	1,193	370	1,000	210	942	180	882	145	800	120
May-December	1,883	1,270	1,648	930	1,413	600	1,179	340	985	180	914	170	850	135	675	75	564	45
June	1,750	1,150	1,333	500	982	180	890	150	816	130	589	55	530	30	450	11	350	4
July	860	140	600	60	566	45	531	30	492	20	416	7	343	4	284	2	242	1
August-October	900	160	511	25	443	10	427	8	296	3	277	2	258	1	239	1	210	0
November	1,632	910	1,106	275	920	175	825	130	610	65	563	45	513	25	455	12	343	4

PRELIMINARY DRAFT

* Percent of time habitat unit (HU) is equalled or exceeded at the given flow (cfs).

**Appendix J: Habitat Units Lost from Inundation in the Little Cypress Bayou. The Value of
Variables Used to Calculate the HSI Were Determined from the Hwy 154 Downstream
Transect According to Table G1**

Species	Month Discharge (cfs)* /Acres**																							
	January 149/211		February 253/305		March 298/346		April 206/262		May 193/252		June 45/114		July 6/79		August 2/76		September 2/76		October 3/76		November 33/104		December 92/158	
	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU	HSI	HU
Spotted bass	0.79	167	0.79	241	0.65	225	0.76	199	0.77	149	0.67	76	0.62	49	0.62	47	0.62	47	0.62	47	0.67	70	0.79	125
Spotted sucker	0.49	103	0.60	163	0.49	169	0.48	126	0.43	83	0.40	46	0.32	25	0.36	27	0.36	27	0.36	27	0.40	42	0.42	66
Brook silverside	0.43	91	0.37	113	0.34	117	0.41	107	0.41	79	0.45	51	0.38	30	0.38	29	0.38	29	0.38	29	0.41	43	0.45	71
Blacktail shiner	0.79	167	0.90	274	0.90	311	0.89	233	0.82	158	0.56	64	0.52	41	0.52	39	0.52	39	0.52	39	0.56	58	0.73	115
Ironcolor shiner	0.49	103	0.40	122	0.37	128	0.42	110	0.42	81	0.37	42	0.36	28	0.45	34	0.45	34	0.45	34	0.37	38	0.37	58
Longear sunfish	0.48	101	0.42	128	0.34	117	0.38	99	0.41	79	0.43	49	0.43	34	0.49	37	0.49	37	0.49	37	0.43	45	0.49	77
Pickereel	0.45	95	0.43	131	0.42	145	0.45	118	0.43	83	0.62	71	0.60	47	0.60	46	0.60	46	0.60	46	0.62	64	0.65	103
Flathead catfish	0.42	89	0.45	137	0.46	159	0.43	113	0.42	81	0.32	36	0.31	24	0.39	30	0.39	30	0.39	30	0.33	34	0.36	57
Slough darter	0.45	95	0.39	119	0.38	131	0.41	107	0.43	83	0.46	52	0.46	36	0.58	44	0.58	44	0.58	44	0.46	48	0.56	88
Total	--	1,011	--	1,448	--	1,502	--	1,212	--	876	--	487	--	314	--	333	--	333	--	333	--	333	--	760

PRELIMINARY DRAFT